

Reduced basis for variational inequalities Application to contact mechanics

Amina Benaceur^{1,2,3}, V. Ehrlacher¹ and A. Ern¹

¹Massachusetts Institute of Technology, USA

²University Paris-Est, CERMICS (ENPC) and INRIA Paris, France

³EDF R&Dg, France



February 19th, 2020



Motivations for contact


- Long-standing question in many contexts...
- Present industrial context
 - Behavior of valve components
 - Costly simulations ($\approx 12\text{h}$ using 'code_aster')
- Nonlinearities, constraints

Challenge

- Expensive for parameterized studies

Goal

- Nonlinearly constrained model reduction



Variational inequalities with linear constraints

Reduced basis procedures

-  [*Haasdonk, Salomon, Wohlmuth.*](#) A reduced basis method for parametrized variational inequalities. [2012](#).
-  [*Balajewicz, Amsallem, Farhat.*](#) Projection-based model reduction for contact problems. [2016](#).
-  [*Fauque, Ramière, Ryckelynck.*](#) Hybrid hyper-reduced modeling for contact mechanics problems. [2018](#).

Related work

-  [*Burkovska, Haasdonk, Salomon, Wohlmuth.*](#) Reduced basis methods for pricing options with the Black-Scholes and Heston models. [2014](#).
-  [*Glas, Urban.*](#) Numerical investigations of an error bound for reduced basis approximations of noncoercive variational inequalities. [2015](#).
-  [*Bader, Zhang, Veroy.*](#) An empirical interpolation approach to reduced basis approximations for variational inequalities, [2016](#).
-  [*Burkovska.*](#) PhD thesis. [2016](#).

Outline

① Elastic contact problem

② Abstract model problem

③ The reduced-basis model

Elastic contact problem

Strain tensor

$$\varepsilon(v) := \frac{1}{2}(\nabla v + \nabla v^T)$$

Stress tensor

$$\sigma(v) = \frac{E\nu}{(1+\nu)(1+2\nu)} \text{tr}(\varepsilon(v))\mathcal{I} + \frac{E}{(1+\nu)}\varepsilon(v)$$

E : Young modulus ν : Poisson coefficient

Parametric equilibrium condition


$$\nabla \cdot \sigma(u(\mu)) = \ell(\mu) \quad \text{in } \Omega(\mu)$$

$$a(\mu; v, w) = \int_{\Omega(\mu)} \sigma(v) : \varepsilon(w) \quad \text{and} \quad f(\mu; w) = \int_{\Omega(\mu)} \ell(\mu) w$$


Other nonlinearities can be handled

Non-interpenetration condition

Initial configuration Deformed configuration



Non-interpenetration condition



Consider $\mathcal{V}(\mu) = H^1(\Omega_1(\mu)) \times H^1(\Omega_2(\mu))$


Admissible solutions are denoted $u(\mu) = (u_1(\mu), u_2(\mu)) \in \mathcal{V}(\mu)$

For all $z \in \Gamma_1^c(\mu)$

$$\underbrace{(u_1(\mu)(z) - (u_2(\mu) \circ \rho(\mu, u(\mu))) (z))}_{\text{displacement difference}} \cdot \tilde{n}_2(\mu, u(\mu))(z) \geq \underbrace{(\rho(\mu, u(\mu))(z) - z)}_{\text{initial gap}} \cdot \tilde{n}_2(\mu, u(\mu))(z)$$

\implies Proof: Benaceur, PhD, 2018.

Non-interpenetration condition



Consider $\mathcal{V}(\mu) = H^1(\Omega_1(\mu)) \times H^1(\Omega_2(\mu))$

Admissible solutions are denoted $u(\mu) = (u_1(\mu), u_2(\mu)) \in \mathcal{V}(\mu)$

For all $z \in \Gamma_1^c(\mu)$

$$(u_1(\mu)(z) - \underbrace{(u_2(\mu) \circ \rho(\mu, u(\mu))) (z)}_{-k(\mu, \mathbf{u}(\mu); \mathbf{u}(\mu))}) \cdot \underbrace{\tilde{n}_2(\mu, u(\mu))(z)}_{\tilde{n}_2(\mu, u(\mu))(z)} \geq \underbrace{(\rho(\mu, u(\mu))(z) - z)}_{-g(\mu, \mathbf{u}(\mu))} \cdot \underbrace{\tilde{n}_2(\mu, u(\mu))(z)}_{\tilde{n}_2(\mu, u(\mu))(z)}$$

\implies Proof: Benaceur, PhD, 2018.

Model problem

$\Omega(\mu)$: bounded domain in \mathbb{R}^d with a contact boundary $\Gamma_1^c(\mu) \subset \partial\Omega(\mu)$

$\mathcal{V}(\mu)$: Hilbert space on $\Omega(\mu)$

\mathcal{P} : parameter set

For many values $\mu \in \mathcal{P}$: Find $u(\mu) \in \mathcal{V}$ such that

$$\begin{aligned} u(\mu) &= \underset{v \in \mathcal{V}(\mu)}{\operatorname{argmin}} \frac{1}{2} a(\mu; v, v) - f(\mu; v) \\ k(\mu, \textcolor{blue}{u}(\mu); \textcolor{green}{u}(\mu)) &\leq g(\mu, \textcolor{blue}{u}(\mu)) \quad \text{a.e. on } \Gamma_1^c(\mu) \end{aligned}$$

$k(\mu, \cdot; \cdot)$ is semi-linear : natural for contact problems
handy for iterative solution methods

Lagrangian formulation

We consider

- the convex cone $\mathcal{W}(\mu) := L^2(\Gamma_1^c(\mu), \mathbb{R}_+)$
- the Lagrangian $\mathcal{L}(\mu) : \mathcal{V}(\mu) \times \mathcal{W}(\mu) \rightarrow \mathbb{R}$ defined as

$$\mathcal{L}(\mu)(\mathbf{v}, \boldsymbol{\eta}) := \frac{1}{2}a(\mu; \mathbf{v}, \mathbf{v}) - f(\mu; \mathbf{v}) + \left(\int_{\Gamma_1^c(\mu)} k(\mu, \mathbf{v}; \mathbf{v}) \boldsymbol{\eta} - \int_{\Gamma_1^c(\mu)} g(\mu, \mathbf{v}) \boldsymbol{\eta} \right)$$

Find $(\mathbf{u}(\mu), \lambda(\mu)) \in \mathcal{V}(\mu) \times \mathcal{W}(\mu)$ such that

$$(\mathbf{u}(\mu), \lambda(\mu)) = \arg \min_{\mathbf{v} \in \mathcal{V}(\mu)} \max_{\boldsymbol{\eta} \in \mathcal{W}(\mu)} \mathcal{L}(\mu)(\mathbf{v}, \boldsymbol{\eta})$$

$\mathbf{u}(\mu)$ is the primal solution, $\lambda(\mu)$ is the dual solution

Discrete FEM formulation

- ① FEM space ($\mathcal{N} \gg 1$)

$$V_{\mathcal{N}}(\mu) := \text{span}\{\phi_1(\mu), \dots, \phi_{\mathcal{N}}(\mu)\} \subset \mathcal{V}(\mu)$$

- ② FEM convex cone ($\mathcal{R} \gg 1$)

$$W_{\mathcal{R}}(\mu) := \text{span}_+ \{\psi_1(\mu), \dots, \psi_{\mathcal{R}}(\mu)\} \subset \mathcal{W}(\mu)$$

Discrete FEM formulation

- ① FEM space ($\mathcal{N} \gg 1$)

$$V_{\mathcal{N}}(\mu) := \text{span}\{\phi_1(\mu), \dots, \phi_{\mathcal{N}}(\mu)\} \subset \mathcal{V}(\mu)$$

- ② FEM convex cone ($\mathcal{R} \gg 1$)

$$W_{\mathcal{R}}(\mu) := \text{span}_+ \{\psi_1(\mu), \dots, \psi_{\mathcal{R}}(\mu)\} \subset \mathcal{W}(\mu)$$

Algebraic FEM formulation

$$(\mathbf{u}(\mu), \boldsymbol{\lambda}(\mu)) = \arg \min_{\mathbf{v} \in \mathbb{R}^{\mathcal{N}}, \boldsymbol{\eta} \in \mathbb{R}_{+}^{\mathcal{R}}} \left\{ \frac{1}{2} \mathbf{v}^T \mathbf{A}(\mu) \mathbf{v} - \mathbf{v}^T \mathbf{f}(\mu) + \boldsymbol{\eta}^T (\mathbf{K}(\mu, \mathbf{v}) \mathbf{v} - \mathbf{g}(\mu, \mathbf{v})) \right\}$$

$$\mathbf{A}(\mu)_{ij} = a(\mu; \phi_j(\mu), \phi_i(\mu)) \quad \mathbf{K}(\mu, w)_{ij} = \int_{\Gamma_1^c(\mu)} k(\mu, w; \phi_j(\mu)) \psi_i(\mu)$$

Additional nonlinearity caused by spatial discretization

The Kačanov method

- Iterative procedure
- Consists in solving the following problems : For all $k \geq 1$

$$(\mathbf{u}^k(\mu), \boldsymbol{\lambda}^k(\mu)) = \arg \min_{\mathbf{v} \in \mathbb{R}^N, \boldsymbol{\eta} \in \mathbb{R}_+^{\mathcal{R}}} \frac{1}{2} \mathbf{v}^T \mathbf{A}(\mu) \mathbf{v} - \mathbf{v}^T \mathbf{f}(\mu)$$

$$+ \boldsymbol{\eta}^T (\mathbf{K}(\mu, \mathbf{u}^{k-1}(\mu)) \mathbf{v} - \mathbf{g}(\mu, \boldsymbol{\lambda}^{k-1}(\mu)))$$

- Stopping criteria

$$\frac{\|\mathbf{u}^k(\mu) - \mathbf{u}^{k-1}(\mu)\|_{\mathbb{R}^N}}{\|\mathbf{u}^{k-1}(\mu)\|_{\mathbb{R}^N}} \leq \epsilon_{\text{ka}}^{\text{pr}} \quad \frac{\|\boldsymbol{\lambda}^k(\mu) - \boldsymbol{\lambda}^{k-1}(\mu)\|_{\mathbb{R}^N}}{\|\boldsymbol{\lambda}^{k-1}(\mu)\|_{\mathbb{R}^N}} \leq \epsilon_{\text{ka}}^{\text{du}}$$

The reduced-basis model: Reference configuration

Geometric mapping $h(\mu)$ defined on a reference domain $\breve{\Omega}$ (with $I = 2$)

$$h(\mu) : \breve{\Omega} \rightarrow \Omega(\mu)$$

$$x \mapsto \sum_{i=1}^I h_i(\mu, x) \mathbf{1}_{\breve{\Omega}_i}(x)$$

The reduced-basis model: Reference configuration

Geometric mapping $h(\mu)$ defined on a reference domain $\breve{\Omega}$ (with $I = 2$)

$$h(\mu) : \breve{\Omega} \rightarrow \Omega(\mu)$$

$$x \mapsto \sum_{i=1}^I h_i(\mu, x) \mathbf{1}_{\breve{\Omega}_i}(x)$$

$$\Omega_i(\mu) = h(\mu)(\breve{\Omega}_i) \quad \forall i \in \{1, 2\}$$

$$\Gamma_i^c(\mu) = h(\mu)(\breve{\Gamma}_i^c) \quad \forall i \in \{1, 2\}$$

Reference Hilbert space

$$\check{\mathcal{V}} := H^1(\breve{\Omega}; \mathbb{R}^d)$$

Parametric Hilbert space

$$\mathcal{V}(\mu) = \check{\mathcal{V}} \circ h(\mu)^{-1}$$

Reference convex cone

$$\check{\mathcal{W}} := L^2(\breve{\Gamma}_1^c; \mathbb{R}_+)$$

Parametric convex cone

$$\mathcal{W}(\mu) = \check{\mathcal{W}} \circ h(\mu)^{-1}|_{\Gamma_1^c(\mu)}$$

Reduced basis spaces

Primal RB subspace

$$\check{V}_N \subset \check{V}_{\mathcal{N}} \subset \check{\mathcal{V}}$$

$$\check{V}_N = \text{span}\{\check{\theta}_1, \dots, \check{\theta}_N\}$$

Dual RB subcone

$$\check{W}_R \subset \check{W}_{\mathcal{R}} \subset \check{\mathcal{W}}$$

$$\check{W}_R = \text{span}_+ \{\check{\xi}_1, \dots, \check{\xi}_R\}$$

Reduced basis spaces

Primal RB subspace

$$\check{V}_N \subset \breve{V}_N \subset \check{\mathcal{V}}$$

$$\check{V}_N = \text{span}\{\check{\theta}_1, \dots, \check{\theta}_N\}$$

Dual RB subcone

$$\check{W}_R \subset \breve{W}_{\mathcal{R}} \subset \check{\mathcal{W}}$$

$$\check{W}_R = \text{span}_+ \{\check{\xi}_1, \dots, \check{\xi}_R\}$$

RB approximations

$$\hat{u}(\mu) = \sum_{n=1}^N \hat{u}_n(\mu) \check{\theta}_n \circ h(\mu)^{-1} \quad \hat{\lambda}(\mu) = \sum_{n=1}^R \hat{\lambda}_n(\mu) \check{\xi}_n \circ h(\mu)^{-1}$$

Reduced problem

$$(\hat{\mathbf{u}}(\mu), \hat{\boldsymbol{\lambda}}(\mu)) = \arg \min_{\hat{\mathbf{v}} \in \mathbb{R}^N, \hat{\boldsymbol{\eta}} \in \mathbb{R}_+^R} \left\{ \frac{1}{2} \hat{\mathbf{v}}^T \hat{\mathbf{A}}(\mu) \hat{\mathbf{v}} - \hat{\mathbf{v}}^T \hat{\mathbf{f}}(\mu) + \hat{\boldsymbol{\eta}}^T (\hat{\mathbf{K}}(\mu, \hat{\mathbf{v}}) \hat{\mathbf{v}} - \hat{\mathbf{g}}(\mu, \hat{\mathbf{v}})) \right\}$$

$$\hat{\mathbf{A}}(\mu) \in \mathbb{R}^{N \times N} \quad \hat{\mathbf{f}}(\mu) \in \mathbb{R}^N \quad \hat{\mathbf{K}}(\mu, \hat{\mathbf{v}}) \in \mathbb{R}^{R \times N} \quad \hat{\mathbf{g}}(\mu, \hat{\mathbf{v}}) \in \mathbb{R}^R$$

Reduced problem

$$(\hat{\mathbf{u}}(\mu), \hat{\boldsymbol{\lambda}}(\mu)) = \arg \min_{\hat{\mathbf{v}} \in \mathbb{R}^N, \hat{\boldsymbol{\eta}} \in \mathbb{R}_+^R} \left\{ \frac{1}{2} \hat{\mathbf{v}}^T \hat{\mathbf{A}}(\mu) \hat{\mathbf{v}} - \hat{\mathbf{v}}^T \hat{\mathbf{f}}(\mu) + \hat{\boldsymbol{\eta}}^T (\hat{\mathbf{K}}(\mu, \hat{\mathbf{v}}) \hat{\mathbf{v}} - \hat{\mathbf{g}}(\mu, \hat{\mathbf{v}})) \right\}$$

$$\hat{\mathbf{A}}(\mu) \in \mathbb{R}^{N \times N} \quad \hat{\mathbf{f}}(\mu) \in \mathbb{R}^N \quad \hat{\mathbf{K}}(\mu, \hat{\mathbf{v}}) \in \mathbb{R}^{R \times N} \quad \hat{\mathbf{g}}(\mu, \hat{\mathbf{v}}) \in \mathbb{R}^R$$

$$\hat{\mathbf{A}}(\mu)_{pn} = a(\mu; \check{\theta}_n \circ h(\mu)^{-1}, \check{\theta}_p \circ h(\mu)^{-1})$$

$$\hat{\mathbf{K}}(\mu, \hat{\mathbf{v}})_{pn} = \int_{\Gamma_1^c(\mu)} k \left(\mu, \sum_{i=1}^N \hat{v}_i \check{\theta}_i \circ h(\mu)^{-1}; \check{\theta}_n \circ h(\mu)^{-1} \right) \check{\xi}_p \circ h(\mu)^{-1}$$

Small (dense) matrices but require FEM reconstructions $\theta_n \circ h(\mu)^{-1}$...etc
 \implies Need to separate (n, p) - and μ -dependencies

Functional separation

For the stiffness matrix, we need a decomposition of the form

$$\left(\hat{\mathbf{A}}(\mu)\right)_{np} := \hat{\mathbf{A}}(\mu, n, p) = \sum_{j=1}^{M^a} \alpha_j(\mu) \hat{\mathbf{A}}_{j,np}$$

- matrices $\{\hat{\mathbf{A}}_j\}_{j=1}^{M^a}$ are **precomputed offline**
- coefficients $\{\alpha_j(\mu)\}_{j=1}^{M^a}$ are **evaluated online**
- straightforward for affine transformations

For the matrix $\hat{\mathbf{K}}(\mu, \hat{\mathbf{v}})$, search for a **separated approximation**

$$k(\mu, u(\mu); \phi_n)(h(\mu)(\check{x})) =: \kappa(\mu, n, \check{x}) \approx \sum_{j=1}^{M^k} \varphi_j(\mu) q_j(n, \check{x})$$

The functions $\{\varphi_j, q_j\}_{j=1}^{M^k}$ are built **offline using the EIM**
 [Barrault, Maday, Nguyen, Patera \('04\)](#)

Offline/Online efficient RB problem

Offline : using $\{q_j\}_{j=1}^{M^k}$, we build

- a matrix $\mathbf{B} \in \mathbb{R}^{M^k \times M^k}$
- a family of matrices $\{\mathbf{C}_j\}_{j=1}^{M^k}$ all in $\mathbb{R}^{R \times N}$

Online : For each new parameter $\mu \in \mathcal{P}$ and a vector $\hat{\mathbf{v}} \in \mathbb{R}^N$

- compute a family of functions $\{\hat{\phi}_j(\mu, \hat{\mathbf{v}})\}_{j=1}^{M^k}$
- assemble

$$\hat{\mathbf{K}}(\mu, \hat{\mathbf{v}}) \approx \mathbf{D}^\kappa(\mu, \hat{\mathbf{v}}) := \sum_{i,j=1}^{M^k} \mathbf{C}_j \mathbf{B}_{ji} \hat{\phi}_i(\mu, \hat{\mathbf{v}}) \in \mathbb{R}^{R \times N}$$

Similarly, we build an efficient approximation $\hat{\gamma}(\mu, \hat{\mathbf{v}})$ for the gap function

Offline/Online efficient RB problem

Solve

$$(\hat{\mathbf{u}}(\mu), \hat{\boldsymbol{\lambda}}(\mu)) = \arg \min_{\hat{\mathbf{v}} \in \mathbb{R}^N, \hat{\boldsymbol{\eta}} \in \mathbb{R}_+^R} \left\{ \frac{1}{2} \hat{\mathbf{v}}^T \hat{\mathbf{A}}(\mu) \hat{\mathbf{v}} - \hat{\mathbf{v}}^T \hat{\mathbf{f}}(\mu) + \hat{\boldsymbol{\eta}}^T (\mathbf{D}^\kappa(\mu, \hat{\mathbf{v}}) \hat{\mathbf{v}} - \mathbf{D}^\gamma \hat{\gamma}(\mu, \hat{\mathbf{v}})) \right\}$$

$$\mathbf{D}^\kappa(\mu, \hat{\mathbf{v}}) \in \mathbb{R}^{R \times N} \quad \mathbf{D}^\gamma \in \mathbb{R}^{R \times M^\gamma} \quad \hat{\gamma}(\mu, \hat{\mathbf{v}}) \in \mathbb{R}^{M^\gamma}$$

- $\mathbf{D}^\kappa(\mu, \hat{\mathbf{v}})$ results from the EIM on κ
- \mathbf{D}^γ results from the EIM on γ

Online stage

Algorithm 1 Online stage

Input : μ

$$\begin{aligned} & \{\hat{\mathbf{f}}_j\}_{1 \leq j \leq J^f}, \{\hat{\mathbf{A}}_j\}_{1 \leq j \leq J^a} \\ & \{(n_i^\kappa, x_i^\kappa)\}_{1 \leq i \leq M^k}, \{q_j^\kappa\}_{1 \leq j \leq M^k} \\ & \{x_i^\gamma\}_{1 \leq i \leq M^g}, \{q_j^\gamma\}_{1 \leq j \leq M^g}, \mathbf{B}^\kappa, \{\mathbf{C}_j^\kappa\}_{1 \leq j \leq M^k} \text{ and } \mathbf{D}^\gamma \end{aligned}$$

- 1: Assemble the vector $\hat{\mathbf{f}}(\mu)$ and the matrix $\hat{\mathbf{A}}(\mu)$
- 2: Compute $\hat{\kappa}(\mu, \hat{\mathbf{v}})$ and $\hat{\gamma}(\mu, \hat{\mathbf{v}})$ EIM on the contact map
- 3: Compute $\mathbf{D}^\kappa(\mu)$ using $\hat{\kappa}(\mu, \hat{\mathbf{v}})$ EIM on the gap map
- 4: Solve the reduced saddle-point problem to obtain $\hat{\mathbf{u}}(\mu)$ and $\hat{\lambda}(\mu)$

Output : $\hat{\mathbf{u}}(\mu)$ and $\hat{\lambda}(\mu)$

Basis constructions

Two goals

- ① Build $\check{V}_N \subset \check{V}_{\mathcal{N}}$ of dimension $N \ll \mathcal{N}$
⇒ POD ✓
- ② Build $\check{W}_R \subset \check{W}_{\mathcal{R}}$ of dimension $R \ll \mathcal{R}$
Requirement: Positive basis vectors

Basis constructions

Two goals

- ① Build $\check{V}_N \subset \check{V}_{\mathcal{N}}$ of dimension $N \ll \mathcal{N}$
 \Rightarrow POD ✓

- ② Build $\check{W}_R \subset \check{W}_{\mathcal{R}}$ of dimension $R \ll \mathcal{R}$

Requirement: Positive basis vectors

- POD ✗
- NMF ✓ : considered in *Balajewicz, Amsallem, Farhat ('16)*
- Angle-greedy algorithm ✓ : introduced in *Haasdonk, Salomon, Wohlmuth ('12)*
- Cone-projected greedy algorithm ✓✓ : devised in this work (19')

Non-negative Matrix Factorization

We search for \mathbf{W}

$$\mathbf{W} = \text{NMF}(\mathbf{T}, \mathcal{R})$$

- Input: integer \mathcal{R}
- Output: \mathcal{R} positive vectors



Lee, Seung ('01)

Non-negative Matrix Factorization

We search for \mathbf{W}

$$\mathbf{W} = \text{NMF}(\mathbf{T}, R)$$

- Input: integer R
- Output: R positive vectors



Lee, Seung ('01)

NMF optimization problem

$$(\mathbf{W}, \mathbf{H}) = \underset{\begin{array}{c} \tilde{\mathbf{W}} \in \mathbb{R}_+^{R \times R} \\ \tilde{\mathbf{H}} \in \mathbb{R}_+^{R \times P} \end{array}}{\operatorname{argmin}} \|\mathbf{T} - \tilde{\mathbf{W}}\tilde{\mathbf{H}}\|^2$$

Functional $\|\mathbf{T} - \tilde{\mathbf{W}}\tilde{\mathbf{H}}\|$ is not convex in both variables $\tilde{\mathbf{W}}$ and $\tilde{\mathbf{H}}$ together \implies only local minima

- Non-unique solution

Non-negative Matrix Factorization

We search for \mathbf{W}

$$\mathbf{W} = \text{NMF}(\mathbf{T}, \mathcal{R})$$

- Input: integer \mathcal{R}
- Output: \mathcal{R} positive vectors



Lee, Seung ('01)

NMF optimization problem

$$(\mathbf{W}, \mathbf{H}) = \underset{\begin{array}{c}\tilde{\mathbf{W}} \in \mathbb{R}_+^{\mathcal{R} \times R} \\ \tilde{\mathbf{H}} \in \mathbb{R}_+^{R \times P}\end{array}}{\operatorname{argmin}} \|\mathbf{T} - \tilde{\mathbf{W}} \mathbf{Z} \mathbf{Z}^{-1} \tilde{\mathbf{H}}\|^2$$

Functional $\|\mathbf{T} - \tilde{\mathbf{W}} \tilde{\mathbf{H}}\|$ is not convex in both variables $\tilde{\mathbf{W}}$ and $\tilde{\mathbf{H}}$ together \implies only local minima

- Non-unique solution

Non-negative Matrix Factorization

We search for \mathbf{W}

$$\mathbf{W} = \text{NMF}(\mathbf{T}, R)$$

- Input: integer R 😞
- Output: R positive vectors
 - ❑ Lee, Seung('01)

NMF optimization problem

$$(\tilde{\mathbf{W}}, \tilde{\mathbf{H}}) = \underset{\substack{\tilde{\mathbf{W}} \in \mathbb{R}_+^{R \times R} \\ \tilde{\mathbf{H}} \in \mathbb{R}_+^{R \times P}}}{\operatorname{argmin}} \| \mathbf{T} - \tilde{\mathbf{W}} \mathbf{Z} \mathbf{Z}^{-1} \tilde{\mathbf{H}} \|^2$$

Functional $\| \mathbf{T} - \tilde{\mathbf{W}} \tilde{\mathbf{H}} \|$ is not convex in both variables $\tilde{\mathbf{W}}$ and $\tilde{\mathbf{H}}$ together \implies only local minima

- Non-unique solution 😞

Angle-greedy algorithm

Selection criterion

$$\mu_n \in \operatorname{argmax}_{\mu \in \mathcal{P}^{\text{tr}}} \| \nabla \left(\check{\lambda}(\mu; \cdot), \check{W}_{n-1} \right) \|_{\ell^\infty(\check{\Gamma}_1^{c, \text{tr}})}$$

Dual set at iteration n

$$\check{K}_n := \operatorname{span}_+ \{ \check{\lambda}(\mu_1; \cdot), \dots, \check{\lambda}(\mu_n; \cdot) \}$$

Stopping criterion

$$r_n < \epsilon_{\text{du}}$$

- Input: Tolerance ϵ_{du} to reach
- Output: R positive vectors
- \check{W}_n is a linear space

Angle-greedy algorithm

Selection criterion

$$\mu_n \in \operatorname{argmax}_{\mu \in \mathcal{P}^{\text{tr}}} \|\check{\lambda}(\mu; \cdot) - \Pi_{\widetilde{W}_{n-1}}(\check{\lambda}(\mu; \cdot))\|_{\ell^\infty(\check{\Gamma}_1^{c,\text{tr}})}$$

Dual set at iteration n

$$\check{K}_n := \operatorname{span}_+ \{\check{\lambda}(\mu_1; \cdot), \dots, \check{\lambda}(\mu_n; \cdot)\}$$

Stopping criterion

$$r_n < \epsilon_{\text{du}}$$

- Input: Tolerance ϵ_{du} to reach
- Output: R positive vectors
- \widetilde{W}_n is a linear space

Angle-greedy algorithm

Selection criterion

$$\mu_n \in \operatorname{argmax}_{\mu \in \mathcal{P}^{\text{tr}}} \|\check{\lambda}(\mu; \cdot) - \Pi_{\widetilde{W}_{n-1}}(\check{\lambda}(\mu; \cdot))\|_{\ell^\infty(\check{\Gamma}_1^{c,\text{tr}})}$$

Dual set at iteration n

$$\check{K}_n := \operatorname{span}_+ \{\check{\lambda}(\mu_1; \cdot), \dots, \check{\lambda}(\mu_n; \cdot)\}$$

Stopping criterion

$$r_n < \epsilon_{\text{du}}$$

- Input: Tolerance ϵ_{du} to reach 😊
- Output: R positive vectors
- \widetilde{W}_n is a linear space 😞

Cone-projected greedy algorithm

Selection criterion

$$\mu_n \in \operatorname{argmax}_{\mu \in \mathcal{P}^{\text{tr}}} \|\check{\lambda}(\mu; \cdot) - \Pi_{\check{K}_{n-1}}(\check{\lambda}(\mu; \cdot))\|_{\ell^\infty(\check{\Gamma}_1^{c,\text{tr}})}$$

Dual set at iteration n

$$\check{K}_n := \operatorname{span}_+ \{\check{K}_{n-1}, \check{\lambda}(\mu_n; \cdot)\}$$

Stopping criterion

$$r_n < \epsilon_{\text{du}}$$

→ Python cvxopt library for positive projections

- Input: Tolerance ϵ_{du} to reach 😊
- Output: R positive vectors

Cone-projected greedy algorithm

Algorithm 2 Cone-projected weak greedy algorithm

Input : \mathcal{P}^{tr} , $\check{\Gamma}_1^c$ and $\epsilon_{\text{du}} > 0$

- 1: Compute $\mathcal{S}_{\text{du}} = \{\check{\lambda}(\mu; \cdot)\}_{\mu \in \mathcal{P}^{\text{tr}}}$
- 2: Set $\check{K}_0 = \{0\}$, $n = 1$ and $r_1 = \max_{\mu \in \mathcal{P}^{\text{tr}}} \|\check{\lambda}(\mu; \cdot)\|_{\ell^\infty(\check{\Gamma}_1^c)}$
- 3: **while** ($r_n > \epsilon_{\text{du}}$) **do**
- 4: Search $\mu_n \in \operatorname{argmax}_{\mu \in \mathcal{P}^{\text{tr}}} \|\check{\lambda}(\mu; \cdot) - \Pi_{\check{K}_{n-1}}(\check{\lambda}(\mu; \cdot))\|_{\ell^\infty(\check{\Gamma}_1^{c,\text{tr}})}$
- 5: Set $\check{K}_n := \text{span}_+ \{\check{K}_{n-1}, \check{\lambda}(\mu_n; \cdot)\}$
- 6: Set $n = n + 1$
- 7: Set $r_n := \max_{\mu \in \mathcal{P}^{\text{tr}}} \|\check{\lambda}(\mu; \cdot) - \Pi_{\check{K}_{n-1}}(\check{\lambda}(\mu; \cdot))\|_{\ell^\infty(\check{\Gamma}_1^c)}$
- 8: **end while**
- 9: Set $R := n - 1$


Output : $\check{W}_R := \check{K}_R$

→ Python cvxopt library for positive projections


Numerical results

Hertz's half-disks problem


- Imposed displacement on the upper half-sphere
- Parametric radius μ for the lower half-sphere
- $\mathcal{P} = [0.9, 1.12]$, $\mathcal{P}^{\text{tr}} = \{0.905 + 0.01i \mid 0 \leq i \leq 22\}$
- Non-matching meshes, $\mathcal{N} = 1350$ nodes, $\mathcal{R} = 51$



Reference configuration







$$\mu = 0.9$$



$$\mu = 1.12$$


High-fidelity solution procedure

- Conforming FEM for displacement, dG \mathbb{P}_0 for contact pressures
 - Collocation at contact nodes (possible instability)
 - Local Averaged Contact (LAC) : macro-elements (two consecutive segments) (Drouet, Hild ('17))




$\mu = 0.9$ (left) / 1.12 (right), coarse (top) / fine (bottom) meshes

Basis constructions



Primal POD basis





Dual basis

Dual basis dimension R as a function of the truncation threshold ϵ_{du}



ϵ_{du}		$5 \cdot 10^{-2}$	10^{-2}	$5 \cdot 10^{-3}$	10^{-3}
NMF	R	4	10	19	20
Cone-projected greedy	R	2	5	6	13

Empirical interpolation

EIM error as a function of the rank M of the EIM approximation.




Error estimation


Error on the minimum energy $e_{ener}(\mu)$ Relative H^1 -error for the displacement field $e_{displ}(\mu)$

Constraint violation

Maximum interpenetration




CPG/NMF(18)



CPG/NMF(5)


Ring on block

- Imposed displacement at the ring's top ends
- Parametric ring's radius $\mu \in \mathcal{P} := [0.95, 1.15]$, $\text{card}(\mathcal{P}^{\text{tr}}) = 21$
- Non-matching meshes, $\mathcal{N} = 2 \times 1590$ dofs, $\mathcal{R} = 50$ dofs
- Reference (left) and deformed (right) configuration




Ring on block

Basis construction



Primal basis



Dual basis


Conclusions and perspectives

Contributions

- ✓ RBM in a general setting
 - varying normals
 - non-matching meshes
- ✓ Cone-projected greedy algorithm
 - outperforms the NMF
 - outperforms the angle-greedy

Prospectives

- ? Extensions to other types of contact
 - friction
 - multibody
 - dynamic
- ? Convergence rates of the CPG



A reduced basis method for parametrized variational inequalities applied to contact mechanics. AB, V. Ehrlacher and A. Ern. *IJNME*, 2019.

